

This schedule is a summary. For further information refer to the online Australian Immunisation Handbook
[The Australian Immunisation Handbook \(health.gov.au\)](https://www.health.gov.au/the-australian-immunisation-handbook)


| | | |
|--------------------------------------|---|--|
| All children, adolescents and adults | Aboriginal children, adolescents and adults | Specified medical conditions - children adolescents and adults |
|--------------------------------------|---|--|

Specified medical conditions funded for Pneumococcal Vaccination [Pneumococcal disease](#)
[List. Risk conditions for pneumococcal disease | The Australian Immunisation Handbook \(health.gov.au\)](#)

- Previous episode of invasive **pneumococcal disease**
- **Functional or anatomical asplenia**, including sickle cell disease or other haemoglobinopathies, splenectomy or hyposplenia
- **Immunocompromising conditions**, including congenital or acquired immune deficiency, haematological malignancies, solid organ transplant, haematopoietic stem cell transplant and HIV infection
- **CSF leak**, including cochlear implants and intracranial shunts
- **Chronic respiratory disease**, including suppurative lung disease, bronchiectasis, cystic fibrosis and chronic lung disease in preterm infants
- **Chronic renal disease**, including relapsing or persistent nephrotic syndrome and chronic renal impairment eGFR <15ml/min (stages 4 and 5) or receiving dialysis

Additional specified medical conditions for children less than 5 years

- Children born **less than 28 weeks gestation**
- **Chronic lung disease** in preterm infants
- **Trisomy 21 (Down Syndrome)**
- **Cardiac disease**, including congenital heart disease, coronary artery disease and heart failure



- ❖ The interval between an age appropriate Prevenar 13 and Pneumovax 23 vaccine should be 12 months. (The minimum interval is 2 months.)
- ❖ Maximum of two Pneumovax 23 vaccines in a lifetime