

All people	Aboriginal and Torres Strait Islander people	Specified medical risk conditions
------------	--	-----------------------------------

Age	Disease	Brand & Route	Important Information
All pregnant women	Influenza	Age appropriate	Every pregnancy, any trimester
	Diphtheria, tetanus, pertussis	Boostrix® (IM) or Adacel® (IM)	Every pregnancy, from 20 weeks
	Respiratory syncytial virus (RSV)	Abrysvo® (IM)	From 28 to 36 weeks. How Respiratory syncytial virus (RSV) The Australian Immunisation Handbook

All adults			
From 18 years	COVID-19	See Recommended COVID-19 vaccines doses	
Under 26 years	Human papillomavirus (HPV)	Gardasil® 9 (IM)	Single dose
			People who are immunocompromised Human papillomavirus (HPV) The Australian Immunisation Handbook 3 dose course
Born during or after 1966	Measles, mumps, rubella	M-M-R II® (SC) (IM) Priorix® (SC) (IM)	For those who have not received two MMR vaccines or who are not immune, give 2 doses, 1 month apart
From 65 years of age	Influenza	Fluad® Quad (IM)	Annually
	Herpes zoster (shingles)	Shingrix® (IM)	2 doses 2-6 months apart. Minimum interval 5 years after a previous NIP funded Zostavax® vaccine and 12 months after previous herpes zoster infection
From 70 years of age	Pneumococcal	Prevenar 13® (IM)	Only single dose needed in adult lifetime Refer to NT Pneumococcal Vaccination Schedule

Aboriginal and Torres Strait Islander adults			
6 months to 64 years	Influenza	Age appropriate	Annually
From 65 years	Influenza	Fluad® Quad (IM)	Annually
20-50 years NT funded	Hepatitis B	Engerix B® Adult (IM)	3 doses (0,1 and 6 months) if not previously vaccinated or who do not have immunity through natural infection
50 years & over with no risk conditions	Pneumococcal	Prevenar 13® (IM) Pneumovax 23® (IM)	Only single dose of Prevenar 13 needed in adult lifetime. Maximum of two Pneumovax 23 vaccines in a lifetime at least 5 years between doses Refer to NT Pneumococcal Vaccination Schedule
	Herpes zoster (shingles)	Shingrix® (IM)	2 doses 2-6 months apart

People with specified medical risk conditions	Influenza	Age appropriate	See Australian Immunisation Handbook for eligible groups with specified medical conditions
	Pneumococcal	Prevenar 13® (IM) Pneumovax 23® (IM)	At diagnosis Prevenar 13® followed by Pneumovax 23® 2-12 months later, then 2 nd dose of Pneumovax 23® at least 5 years later Pneumococcal vaccination for people with medical risk factors
	COVID-19	See Recommended COVID-19 vaccine doses	
	Herpes zoster (shingles)	Shingrix® (IM)	2 doses, 1-2 months apart
	Meningococcal ACWY	Nimenrix® (IM)	See Table. Recommendations for vaccination in people with functional or anatomical asplenia The Australian Immunisation Handbook (health.gov.au)
	Meningococcal B	Bexsero® (IM)	
	<i>Haemophilus influenzae</i> type b (Hib)	Act-HIB® (IM)	

Refugees & other humanitarian entrants aged 20 years and over. Vaccination for migrants, refugees and people seeking asylum in Australia			
20 years and over Check previous vaccination history and or serology before offering catch up	Diphtheria, tetanus, pertussis	Boostrix® (IM) or Adacel® (IM)	3 doses 1 month between doses
	Poliomyelitis	IPOL® (SC)	3 doses 1 month between doses
	Measles, mumps, rubella	M-M-R II® (IM) (SC) Priorix® (IM) (SC)	2 doses 1 month between doses
	Hepatitis B	H-B-Vax II® Adult (IM)	3 doses at 0,1 and 6 months
	Varicella	Varivax® (SC)	2 doses 1 month between doses

This schedule is a summary. For more detailed information refer to the [The Australian Immunisation Handbook \(health.gov.au\)](#)