






























The Right Age for Baby Needles

Based on the NT *Childhood Immunisation Schedule*

Protect your children from serious diseases by getting their free baby needles (vaccinations) at the right time

Age		How many needles	Recommended needles to protect against these diseases
BIRTH	In hospital 		Hepatitis B
6 WEEKS	Baby starting to smile and lift head 		Diphtheria, Tetanus, Pertussis (Whooping Cough), Hepatitis B, Poliomyelitis (Polio), <i>Haemophilus influenzae</i> type b
			Pneumococcal
			Meningococcal B (<i>Aboriginal and some medically at risk children</i>)
		 ORAL	Rotavirus
4 MONTHS	Baby starting to roll over 		Diphtheria, Tetanus, Pertussis (Whooping Cough), Hepatitis B, Poliomyelitis (Polio), <i>Haemophilus influenzae</i> type b
			Pneumococcal
			Meningococcal B (<i>Aboriginal and some medically at risk children</i>)
		 ORAL	Rotavirus
6 MONTHS	Baby starting to sit up 		Diphtheria, Tetanus, Pertussis (Whooping Cough), Hepatitis B, Poliomyelitis (Polio), <i>Haemophilus influenzae</i> type b
			Pneumococcal (<i>Aboriginal and some medically at risk children</i>)
Influenza vaccine – Yearly – from 6 months of age 2 doses – (1 month apart) when having influenza vaccine for the first time. 1 dose each year after that			
12 MONTHS	Baby starting to stand up and walk 		Measles, Mumps, Rubella
			Pneumococcal
			Meningococcal ACWY
			Meningococcal B (<i>Aboriginal and some medically at risk children</i>)
18 MONTHS	Child starting to run 		Measles, Mumps, Rubella and Chicken Pox
			Diphtheria, Tetanus, Pertussis (Whooping Cough)
			<i>Haemophilus influenzae</i> type b
			Hepatitis A (Aboriginal children)
4 YEARS	Before child goes to school 		Diphtheria, Tetanus, Pertussis (Whooping Cough), Poliomyelitis (Polio)
			Hepatitis A (Aboriginal children)
			Pneumococcal (<i>Aboriginal and some medically at risk children</i>)

For more information contact the NT Immunisation Register Information Line on 8922 8315