

### Just **CLICK**, **MIX**, **PREP** to get started.


- ABRYSVO is for **intramuscular injection only**.
- ABRYSVO is for **single use in one patient only**. Discard any residue.
- The lyophilised vaccine (powder) **must be reconstituted only with the diluent provided** using the vial adapter to form ABRYSVO.
- Please review the **Product Information** before administering ABRYSVO.

#### Each ABRYSVO pack contains the following:

Vial containing lyophilised RSVpreF vaccine



Pre-filled syringe containing diluent




Vial adapter



**Note:** needle is not included.

### 1 **CLICK:** Attach vial adapter



- Peel off the top cover from the vial adapter packaging and remove the flip-off cap from the vial.
- While keeping the vial adapter in its packaging, centre over the vial's stopper and **connect with a straight downward push**. Do not push the vial adapter in at an angle as it may result in leaking. Remove the packaging.

### 2

### **MIX:** Reconstitute lyophilised vaccine component to form ABRYSVO



- For all syringe assembly steps, **hold the syringe only by the Luer lock adapter**. This will prevent the Luer lock adapter from detaching during use.
- Twist to remove the syringe cap, then twist to **connect the syringe to the vial adapter**. Stop turning when you feel resistance.
- **Inject the entire contents of the syringe into the vial**. Hold the plunger rod down and gently swirl the vial until the powder is completely dissolved (less than 1 minute). **Do not shake**.

### 3

### **PREP:** Withdraw reconstituted vaccine



- Invert the vial completely and slowly withdraw the **entire contents** into the syringe to **ensure a 0.5 mL dose of ABRYSVO**.
- Twist to **disconnect** the syringe from the vial adapter.
- **Attach a sterile needle** suitable for intramuscular injection.

- The prepared vaccine is a **clear and colourless solution**. Visually inspect the vaccine for large particulate matter and discoloration prior to administration. Do not use if large particulate matter or discoloration is found.
- **ABRYSVO should be administered immediately (within 4 hours) after reconstitution**. Do not store above 30°C. Do not store in a refrigerator (2°C – 8°C). Do not freeze.

Scan to watch a video on how to prepare ABRYSVO (2 mins)



**Reference:** 1. ABRYSVO<sup>®</sup> Product Information.

**Before prescribing, please review the Product Information available at [www.pfizer.com.au](http://www.pfizer.com.au) or by calling 1800 675 229.**

▼ This vaccine is subject to additional monitoring in Australia. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse events at: [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

Alternatively, contact Pfizer Drug Safety at [AUS.AEReporting@pfizer.com](mailto:AUS.AEReporting@pfizer.com). For medical enquiries, please visit [www.pfizermedicalinformation.com.au](http://www.pfizermedicalinformation.com.au) or call 1800 675 229.

**PBS Information:** This product is not listed on the NIP or PBS.