

All people		Aboriginal and Torres Strait Islander people		Specified medical risk conditions		
Age	Disease	Brands & Route	Dose	Important information		
Birth	Hepatitis B	H-B-Vax II® Paed (IM) or Engerix B® Paed (IM)	Birth	Within 7 days of birth (ideally 24 hours). No catch-up required		
6 weeks	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b (Hib)	Infanrix hexa® (IM) or Vaxelis® (IM)	1			
	Pneumococcal	Prevenar 13® (IM)	1			
	Rotavirus	Rotarix® (Oral)	1	Dose 1 must be given before 15 weeks of age		
	Meningococcal B	Bexsero® (IM)	1	Prophylactic paracetamol		
4 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	2			
	Pneumococcal	Prevenar 13® (IM)	2			
	Rotavirus	Rotarix® (Oral)	2	Dose 2 must be given before 25 weeks of age		
	Meningococcal B	Bexsero® (IM)	2	Prophylactic paracetamol		
6 months	I N F L U E N Z A	Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	3	Dose 2 and 3 need to be given 2 months apart	
		Pneumococcal	Prevenar 13® (IM)	3	3 rd dose at 6 months	3 rd dose at 6 months
		Influenza – ANNUALLY In children aged 6 months to less than 9 years, in the first year of administration, give 2 doses, 1 month apart Influenza (flu) The Australian Immunisation Handbook	Age appropriate	Annually	Children aged 6 months to less than 5 years Aboriginal people 6 months and over People 6 months and over with medical risk conditions People with medical conditions that increase their risk of influenza Influenza (flu) AIH	
12 months	I N F L U E N Z A	Measles, mumps, rubella	Priorix® (IM) (SC) or M-M-R II® (IM) (SC)	1	Infants from 6 months of age can receive MMR vaccine for travel to highly endemic areas or during outbreaks. Will still require scheduled doses at 12 and 18 months of age	
		Meningococcal ACWY	Nimenrix® (IM)	1		
		Pneumococcal	Prevenar 13® (IM)	3 or 4		
		Meningococcal B	Bexsero® (IM)	3 or 4	Prophylactic paracetamol Catch-up schedule for children under 2 years till 31 Dec 2026 . If commence course over 12 months of age it is a 2 dose schedule - 8 weeks between doses.	
		Hepatitis B	H-B-Vax II® Paed (IM) or Engerix B® Paed (IM)	4	Only for infants born less than 2000g and/or infants born before 32 weeks gestation (regardless of weight)	
18 months	I N F L U E N Z A	Measles, mumps, rubella, varicella	Priorix-Tetra® (IM) (SC) or ProQuad® (IM) (SC)	2		
		<i>Haemophilus influenzae</i> type b	Act-HIB® (IM)	4		
		Diphtheria, tetanus, pertussis	Infanrix® (IM) or Tripacel® (IM)	4		
		Hepatitis A	VAQTA® (IM)	1		
4 years	I N F L U E N Z A	Diphtheria, tetanus, pertussis, polio	Infanrix IPV® (IM) or Quadracel® (IM)	5		
		Hepatitis A	VAQTA® (IM)	2		
		Pneumococcal	Pneumovax 23	1	See NT Pneumococcal schedule	See NT Pneumococcal schedule
12 years year 7 students	I N F L U E N Z A	Human papillomavirus (HPV)	Gardasil 9® (IM)	1	People who are immunocompromised Human papillomavirus (HPV) AIH	
		Diphtheria, tetanus, pertussis	Boostrix® (IM) or Adacel® (IM)	1		
14 - 19 years	I N F L U E N Z A	Meningococcal ACWY	MenQuadfi® (IM)	2		
		Meningococcal B	Bexsero® (IM)	1 & 2	Catch-up schedule for adolescents 15-19 years till 31 Dec 2026. 8 weeks between doses	
		Pneumococcal	Pneumovax 23® (IM)	2	See NT Pneumococcal schedule	See NT Pneumococcal schedule

This schedule is a summary. For more detailed information refer to the [The Australian Immunisation Handbook \(health.gov.au\)](http://health.gov.au)