

NT Immunisation Schedule

Children and Adolescents up to 19 years

Vaccines funded under the National Immunisation Program/NT Health

October 2026

Ph (08) 8922 8315

Immunisation program | NT Health

KEY

All people

Aboriginal and Torres Strait Islander people

Specified medical risk conditions

Age	Disease	Brands/Route	Dose	Important information	
Birth	Hepatitis B	H-B-Vax II® Paed (IM) Engerix B® Paed (IM)	Birth	Within 7 days of birth (ideally 24 hours). No catch-up required	
6 weeks	Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	1		
	Pneumococcal	Prevenar 20® (IM)	1		
	Rotavirus	Rotarix® (Oral)	1	Dose 1 must be given before 15 weeks of age	
	Meningococcal B	Bexsero® (IM)	1	Prophylactic paracetamol	
4 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	2		
	Pneumococcal	Prevenar 20® (IM)	2		
	Rotavirus	Rotarix® (Oral)	2	Dose 2 must be given before 25 weeks of age	
	Meningococcal B	Bexsero® (IM)	2	Prophylactic paracetamol	
ANNUAL INFLUENZA VACCINE For all children over 6 months to less than 5 years, Aboriginal and Torres Strait Islander children and people with specified medical risk conditions over 6 months to 19 years	6 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	3	Doses 2 and 3, give 2 months apart Doses 1 and 3, give 4 months apart	
		Pneumococcal <i>See NT Pneumococcal Schedule</i>	3	Aboriginal and Torres Strait Islander	Specified medical risk conditions
		Influenza <i>See NT Influenza schedule</i>	Annual	6 months to < 2 years receiving influenza vaccine for first time, 2 doses, 4 weeks apart. <i>See NT Influenza schedule for children with medical risk conditions</i>	
	12 months	Measles, mumps, rubella NT funded	1	Infants from 6 months can receive MMR vaccine for travel overseas or during outbreaks. Will still require scheduled doses at 12 and 18 months	
		Meningococcal ACWY	1		
		Pneumococcal	3 or 4	Aboriginal and Torres Strait Islander	Specified medical risk conditions
		Meningococcal B	3	Prophylactic paracetamol Catch-up for children under 2 years until 31 Dec 2026	
		Hepatitis B	4	Only for infants born less than 2000g and/or infants born before 32 weeks gestation (regardless of weight)	
	18 months	Measles, mumps, rubella, varicella	2		
		<i>Haemophilus influenzae</i> type b	4		
		Diphtheria, tetanus, pertussis	4		
		Hepatitis A	1		
	4 years	Diphtheria, tetanus, pertussis, polio	5		
		Hepatitis A	2		
		Pneumococcal <i>See NT Pneumococcal Schedule</i>		For Aboriginal and Torres Strait Islander children who have not received Prevenar 20®	
	From 12 years or Year 7	Human papillomavirus (HPV)	1	3 doses for medical risk conditions HPV 	
		Diphtheria, tetanus, pertussis	1		
		Pneumococcal <i>See NT Pneumococcal Schedule</i>		For Aboriginal and Torres Strait Islander children who have not received Prevenar 20®	
	14-19 years or Year 9	Meningococcal ACWY	2	Recommended to receive one dose ≥14 years regardless of previous doses	
		Meningococcal B	1 & 2	Recommended to receive 2 doses, 8 weeks apart, ≥14 years regardless of previous doses	

This schedule is a summary. For more detailed information refer to [The Australian Immunisation Handbook](#)