NT Immunisation Schedule

Children and Adolescents up to 19 years

Vaccines funded under the National Immunisation Program/NT Health

November 2026

Ph (08) 8922 8315

Immunisation program | NT Health

KEY: All people | Aboriginal and Torres Strait Islander people | Specified medical risk conditions

Age		Disease	Brands/Route	Dose	Important information
Birth		Hepatitis B	H-B-Vax II®Paed (IM) Engerix B® Paed (IM)	Birth	Within 7 days of birth (ideally 24 hours). No catch-up required
6 weeks		Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	1	
		Pneumococcal	Prevenar 20® (IM)	1	
		Rotavirus	Rotarix® (Oral)	1	Dose 1 must be given before 15 weeks of age
		Meningococcal B	Bexsero® (IM)	1	Prophylactic paracetamol
4 months		Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	2	
		Pneumococcal	Prevenar 20® (IM)	2	
		Rotavirus	Rotarix® (Oral)	2	Dose 2 must be given before 25 weeks of age
		Meningococcal B	Bexsero® (IM)	2	Prophylactic paracetamol
ANNUAL INFLUENZA VACCINE For all children over 6 months to less than 5 years, Aboriginal and Torres Strait Islander children and people with specified medical risk conditions over 6 months to 19 years	**6 months**	Diphtheria, tetanus, pertussis, hepatitis B, polio, Hib	Infanrix hexa® (IM) or Vaxelis® (IM)	3	Doses 2 and 3, give 2 months apart Doses 1 and 3, give 4 months apart
		Pneumococcal ***See NT Pneumococcal Schedule***	Prevenar 20® (IM)	3	Aboriginal and Torres Strait Islander / Specified medical risk conditions
		Influenza ***See NT Influenza schedule***	Vaxigrip® (IM) Flucelvax®(IM)	Annual	**6 months to < 2 years** receiving influenza vaccine for first time, 2 doses, 4 weeks apart. ***See NT Influenza schedule for children with medical risk conditions***
	12 months	Measles, mumps, rubella	Priorix® (IM) (SC)	1	Infants from 6 months can receive MMR vaccine for travel overseas or during outbreaks. Will still require scheduled doses at 12 and 18 months
		Meningococcal ACWY	Nimenrix® (IM)	1	
		Pneumococcal	Prevenar 20® (IM)	3 or 4	Aboriginal and Torres Strait Islander / Specified medical risk conditions
		Meningococcal B	Bexsero® (IM)	3	Prophylactic paracetamol Catch-up for children under 2 years until 31 Dec 2026
		Hepatitis B	H-B-Vax II® Paed (IM) or Engerix B® Paed (IM)	4	Only for infants born less than 2000g and/or infants born before 32 weeks gestation (regardless of weight)
	18 months	Measles, mumps, rubella, varicella	Priorix-Tetra® (IM) (SC)	2	
		Haemophilus influenzae type b	Act-HIB® (IM)	4	
		Diphtheria, tetanus, pertussis	Infanrix® (IM) or Tripacel® (IM)	4	
		Hepatitis A	VAQTA® (IM)	1	
	4 years	Diphtheria, tetanus, pertussis, polio	Infanrix IPV® (IM) or Quadracel® (IM)	5	
		Hepatitis A	VAQTA® (IM)	2	
		Pneumococcal ***See NT Pneumococcal Schedule***	Prevenar 20® (IM)	For Aboriginal and Torres Strait Islander children **up to 18 years of age** who have not received Prevenar 20®	
	From 12 years or Year 7	Human papillomavirus (HPV)	Gardasil 9® (IM)	1	3 doses for medical risk conditions | HPV |
		Diphtheria, tetanus, pertussis	Boostrix® (IM) or Adacel® (IM)	1	
		Pneumococcal ***See NT Pneumococcal Schedule***	Prevenar 20® (IM)	For Aboriginal and Torres Strait Islander children **up to 18 years of age** who have not received Prevenar 20®	
	14-19 years or Year 9	Meningococcal ACWY	MenQuadfi® (IM)	2	Recommended to receive one dose ≥14 years regardless of previous doses
		Meningococcal B	Bexsero® (IM)	1 & 2	Recommended to receive 2 doses, 8 weeks apart, ≥14 years regardless of previous doses

This schedule is a summary. For more detailed information refer to The Australian Immunisation Handbook

NORTHERN TERRITORY GOVERNMENT