

This schedule is a summary. For further information refer to the online Australian Immunisation Handbook  
[The Australian Immunisation Handbook \(health.gov.au\)](https://www.health.gov.au/the-australian-immunisation-handbook)

For COVID-19 Vaccines refer to [Recommendations | COVID-19 | The Australian Immunisation Handbook](#)

All children, adolescents and adults		Aboriginal children, adolescents and adults		Specified medical conditions - children, adolescents and adults	
Eligible groups		Disease	Vaccine brand	Route	Notes
Pregnancy		Influenza	Fluarix Tetra or Vaxigrip Tetra	IM	All pregnant women - any trimester, every pregnancy <i>Refer to Additional Vaccines and Notes</i>
		Diphtheria, tetanus, pertussis	Boostrix or Adacel	IM	From 20 weeks of pregnancy, every pregnancy
All Adults	Under 26 years	Human Papillomavirus (HPV)	Gardasil 9	IM	Single dose <b>Specified medical conditions - <a href="#">HPV and severely immunocompromising conditions</a></b>
	Born during or after 1966	Measles, mumps, rubella (MMR)	M-M-R II or Priorix	SC or IM	<b>For those who have not received two MMR vaccines or who are not immune</b> Give 2 doses, 1 month between doses
	From 65 years of age	Influenza	Fluad Quad	IM	Annually
		COVID-19	Follow <a href="#">Recommendations   COVID-19   The Australian Immunisation Handbook (health.gov.au)</a>		
	From 65 years of age	Herpes Zoster (shingles)	Shingrix	IM	2 doses, 2-6 months between doses. <b>Minimum interval 5 years after a previous NIP funded Zostavax vaccine and 12 months after previous herpes zoster infection</b>
	Non-Aboriginal adults from 70 years	Pneumococcal	Prevenar 13	IM	<i>Refer to NT Pneumococcal Vaccination Schedule and <a href="#">Adults   Pneumococcal disease   The Australian Immunisation Handbook (health.gov.au)</a></i>
	Medically at risk <i>May require extra funded vaccines</i>	Pneumococcal	<a href="#">Pneumococcal vaccination for people with medical risk factors</a>		
		Influenza	<a href="#">List. Specified medical conditions associated with increased risk of influenza disease and severe outcomes   The Australian Immunisation Handbook (health.gov.au)</a>		
		<i>Haemophilus influenzae</i> type b (Hib)	<a href="#">Hib vaccination for people who are immunocompromised</a>		
		Meningococcal ACWY and Meningococcal B	<a href="#">List. Specified medical conditions associated with increased risk of invasive meningococcal disease   The Australian Immunisation Handbook (health.gov.au)</a>		
		COVID-19	<a href="#">People with medical conditions that increase their risk of COVID-19</a>		
		Shingrix	<a href="#">People aged ≥18 years who are immunocompromised or shortly expected to be immunocompromised</a>		
Aboriginal adults	Up to and including 64 years	Influenza	Flurarix Tetra or Vaxigrip Tetra	IM	<a href="#">Influenza vaccination for Aboriginal and Torres Strait Islander people</a>
	From 65 years		Fluad Quad	IM	
	20 to 50 years	Hepatitis B	Engerix B (adult) or H-B-Vax II (adult)	IM	3 doses at 0, 1 and 6 months <b>If not previously vaccinated or who do not have immunity through natural infection</b> Use same vaccine brand to complete course if possible. <a href="#">Hepatitis B vaccination for Aboriginal and Torres Strait people</a>
	50 years and over With no risk conditions	Pneumococcal	Prevenar 13 Pneumovax 23	IM	<i>Refer to NT Pneumococcal Vaccination Schedule and <a href="#">Pneumococcal vaccination for Aboriginal and Torres Strait Islander people</a></i>
		Herpes Zoster (shingles)	Shingrix	IM	2 doses, 2-6 months between doses. <b>Minimum interval 5 years after a previous NIP funded Zostavax vaccine and 12 months after previous herpes zoster infection</b>
<b>Refugees and other humanitarian entrants aged 20 years and over may require funded vaccines</b>  <b>NOTE</b> Check previous vaccination history and/or serology before offering catch up <i>Refer to Australian Immunisation Handbook <a href="#">Vaccination for migrants, refugees and people seeking asylum in Australia</a></i>		Diphtheria, tetanus, pertussis	Boostrix or Adacel	IM	3 doses 1 month between doses
		Poliomyelitis	IPOL	SC	3 doses 1 month between doses
		Measles, mumps, rubella	M-M-R II or Priorix	SC or IM	2 doses 1 month between doses
		Hepatitis B	Engerix B (adult) or H-B-Vax II (adult)	IM	3 doses at 0, 1 and 6 months
		Varicella	Varivax	SC	2 doses 1 month between doses