

This schedule is a summary. For further information refer to the online Australian Immunisation Handbook
[The Australian Immunisation Handbook \(health.gov.au\)](https://www.health.gov.au/the-australian-immunisation-handbook)

For COVID-19 Vaccines refer to [Recommendations | COVID-19 | The Australian Immunisation Handbook](#)

All children, adolescents and adults		Aboriginal children, adolescents and adults		Specified medical conditions - children, adolescents and adults	
Eligible groups		Disease	Vaccine brand	Route	Notes
Pregnancy		Influenza	Fluarix Tetra	IM	All pregnant women - any trimester, every pregnancy <i>Refer to Additional Vaccines and Notes</i>
		Diphtheria, tetanus, pertussis	Boostrix or Adacel	IM	From 20 weeks of pregnancy, every pregnancy
All Adults	Under 26 years	Human Papillomavirus (HPV)	Gardasil 9	IM	Single dose
					Specified medical conditions – Refer to the Australian Immunisation Handbook HPV and severely immunocompromising conditions
	Born after 1966	Measles, mumps, rubella (MMR)	M-M-R II or Priorix	SC or IM	For those who have not received two MMR vaccines or who are not immune Give 2 doses, 1 month between doses
	From 65 years of age	Influenza	Fluad Quad	IM	Annually
		COVID-19	Follow current COVID -19 Australian Immunisation Handbook recommendations		
	From 65 years of age	Herpes Zoster (shingles)	Shingrix	IM	2 doses, 2-6 months between doses. Minimum interval 5 years after a previous NIP funded Zostavax vaccine and 12 months after previous herpes zoster infection Specified medical conditions – refer to the Australian Immunisation Handbook for details People aged ≥18 years who are immunocompromised
					Specified medical conditions – refer to the Australian Immunisation Handbook for details People aged ≥18 years who are immunocompromised
	Non-Aboriginal adults from 70 years	Pneumococcal	Prevenar 13	IM	<i>Refer to NT Pneumococcal Vaccination Schedule and Australian Immunisation Handbook</i>
	Medically at risk <i>May require extra funded vaccines</i>	Pneumococcal	Specified medical conditions - For specific diseases and dosing schedules refer to the Australian Immunisation Handbook		
		Influenza			
		<i>Haemophilus influenzae</i> type b (Hib)	Recommendations for vaccination in people with functional or anatomical asplenia		
		Meningococcal ACWY and Meningococcal B			
COVID-19		Recommendations COVID-19 The Australian Immunisation Handbook (health.gov.au)			
Shingrix		Shingles vaccination schedule from 1 November 2023			
Aboriginal adults	All ages from 6 months	Influenza	As supplied Note age eligibility	IM	<i>Refer to Influenza in Additional Vaccines and Notes</i>
	20 to 50 years	Hepatitis B	Engerix B adult or H-B-Vax II	IM	3 doses (0, 1 and 6 months) If not previously vaccinated or who do not have immunity through natural infection Use same vaccine brand to complete course if possible.
	50 years and over With no risk conditions	Pneumococcal	Prevenar 13 Pneumovax 23	IM	<i>Refer to NT Pneumococcal Vaccination Schedule</i>
		Herpes Zoster (shingles)	Shingrix	IM	2 doses, 2-6 months between doses. Minimum interval 5 years after a previous NIP funded Zostavax vaccine and 12 months after previous herpes zoster infection
Refugees and other humanitarian entrants aged 20 years and over may require funded vaccines NOTE Check previous vaccination history and/or serology before offering catch up <i>Refer to Australian Immunisation Handbook Vaccination for migrants, refugees and people seeking asylum in Australia</i>		Diphtheria, tetanus, pertussis	Boostrix or Adacel	IM	3 doses 1 month between doses
		Poliomyelitis	IPOL	SC	3 doses 1 month between doses
		Measles, mumps, rubella	M-M-R II or Priorix	SC or IM	2 doses 1 month between doses
		Hepatitis B	H-B-Vax II adult	IM	3 doses at 0, 1 and 6 months
		Varicella	Varilrix or Varivax	SC	2 doses 1 month between doses