

This schedule is a summary. For further information refer to the online Australian Immunisation Handbook

[The Australian Immunisation Handbook \(health.gov.au\)](https://www.health.gov.au)

For COVID-19 Vaccines refer to [Recommendations | COVID-19 | The Australian Immunisation Handbook \(health.gov.au\)](https://www.health.gov.au)

All children, adolescents and adults		Aboriginal children, adolescents and adults		Specified medical conditions - children, adolescents and adults	
Age	Disease	Vaccine brand	Dose	Route	Notes
Birth	Hepatitis B	H-B-Vax II (paed) or Engerix-B (paed)	Birth	IM	Administer within 7 days of birth (ideally within 24 hours) no catch up required
6 weeks	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b	Infanrix hexa or Vaxelis	1	IM	ALL CHILDREN
	Pneumococcal	Prevenar 13	1	IM	ALL CHILDREN
	Rotavirus	Rotarix	1	ORAL	Dose 1 must be given before 15 weeks of age
	Meningococcal B	Bexsero	1	IM	ABORIGINAL CHILDREN. Prophylactic paracetamol
4 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b	Infanrix hexa or Vaxelis	2	IM	ALL CHILDREN
	Pneumococcal	Prevenar 13	2	IM	ALL CHILDREN
	Rotavirus	Rotarix	2	ORAL	Dose 2 must be given before 25 weeks of age
	Meningococcal B	Bexsero	2	IM	ABORIGINAL CHILDREN. Prophylactic paracetamol Refer to Additional Vaccines and Notes for children who are immunocompromised
6 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b	Infanrix hexa or Vaxelis	3	IM	ALL CHILDREN. 2 months is required between 2 <sup>nd</sup> and 3 <sup>rd</sup> dose of Infanrix hexa or Vaxelis
	Pneumococcal	Prevenar 13	3	IM	ABORIGINAL CHILDREN receive a 3 <sup>rd</sup> dose at 6 months  Specified medical conditions-receive 3 <sup>rd</sup> dose at 6 months Refer to the NT Pneumococcal Vaccination schedule and AIH <a href="#">Pneumococcal vaccination for people with medical risk factors</a>
6 months to under 5 years	Influenza	Fluarix Tetra or Vaxigrip Tetra	Annually	IM	ALL CHILDREN. If receiving the vaccine for the first time in their life and aged 6 months to under 9 years: give 2 doses, 1 month apart <a href="#">Influenza vaccination 6 months to &lt;5 years</a>
12 months	Measles, mumps, rubella	Priorix or MMR II	1	SC or IM	ALL CHILDREN
	Meningococcal ACWY	Nimenrix	1	IM	ALL CHILDREN
	Pneumococcal	Prevenar 13	3 (or) 4	IM	ALL CHILDREN
	Meningococcal B	Bexsero	3 (or) 4	IM	ABORIGINAL CHILDREN. Prophylactic paracetamol
	Hepatitis B	H-B-Vax II (paed) or Engerix-B (paed)	4	IM	Infants born less than 2000g and/or infants born before 32 weeks gestation (regardless of weight)
18 months	Measles, mumps, rubella, varicella	Priorix-Tetra or ProQuad	2	SC or IM	ALL CHILDREN
	<i>Haemophilus influenzae</i> type b	Act-HIB	4	IM	ALL CHILDREN
	Diphtheria, tetanus, pertussis	Infanrix or Tripacel	4	IM	ALL CHILDREN
	Hepatitis A	VAQTA	1	IM	ABORIGINAL CHILDREN
4 years	Diphtheria, tetanus, pertussis, polio	Infanrix IPV or Quadracel	5	IM	ALL CHILDREN
	Hepatitis A Maximum 2 doses only	VAQTA	2	IM	ABORIGINAL CHILDREN who have NOT had 2 doses previously, 6 months apart
	Pneumococcal	Pneumovax 23	1	IM	ABORIGINAL CHILDREN  Specified medical conditions - refer to the NT Pneumococcal Vaccination schedule and AIH <a href="#">Pneumococcal vaccination for people with medical risk factors</a>
12 years (Year 7 students)	Human Papillomavirus (HPV)	Gardasil 9	1	IM	ALL ADOLESCENTS  Specified medical conditions - 3 dose course: 0, 2 & 6 months schedule Refer to the AIH <a href="#">HPV vaccination for people who are immunocompromised</a>
	Diphtheria, tetanus, pertussis	Boostrix or Adacel	1	IM	ALL ADOLESCENTS
14-19 Years	Meningococcal ACWY	Nimenrix	2	IM	ALL ADOLESCENTS. Vaccine can be given if previous vaccine was given under 14 years
	Pneumococcal	Pneumovax 23 (Dose 2 if not already given)	2	IM	ABORIGINAL ADOLESCENTS  Specified medical conditions - refer to the NT Pneumococcal Vaccination schedule and AIH <a href="#">Pneumococcal vaccination for people with medical risk factors</a>