

All people		Aboriginal and Torres Strait Islander people		Specified medical risk conditions	
Age	Disease	Brands & Route	Dose	Important information	
Birth	Hepatitis B	H-B-Vax II® Paed (IM) or Enderix B® Paed (IM)	Birth	Within 7 days of birth (ideally 24 hours) no catch-up required	
6 weeks	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b	Infanrix hexa® (IM) or Vaxelis® (IM)	1	Dose 1 must be given before 15 weeks of age Prophylactic paracetamol	
	Pneumococcal	Prevenar 13® (IM)	1		
	Rotavirus	Rotarix® (Oral)	1		
	Meningococcal B	Bexsero® (IM)	1		
4 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b	Infanrix hexa® (IM) or Vaxelis® (IM)	2	Dose 2 must be given before 25 weeks of age Prophylactic paracetamol	
	Pneumococcal	Prevenar 13® (IM)	2		
	Rotavirus	Rotarix® (Oral)	2		
	Meningococcal B	Bexsero® (IM)	2		
6 months	Diphtheria, tetanus, pertussis, hepatitis B, polio, <i>Haemophilus influenzae</i> type b	Infanrix hexa® (IM) or Vaxelis® (IM)	3	Dose 2 and 3 need to be given 2 months apart.	
	Pneumococcal	Prevenar 13® (IM)	3	3 rd dose at 6 months	3 rd dose at 6 months
6 months	Influenza	Fluarix® Tetra (IM) or Vaxigrip® Tetra (IM)	Annually	6 months to less than 5 years	If receiving the vaccine for the first time in their life aged over 6 months to less than 9 years, give 2 doses, 1 month apart
				6 months & over	
				6 months & over	
12 months	Measles, mumps, rubella	Priorix® (IM) (SC) or M-M-R II® (IM) (SC)	1	Infants from 6 months of age can receive MMR vaccine for travel to highly endemic areas or during outbreaks. Will still require additional scheduled doses at 12 and 18 months of age	
	Meningococcal ACWY	Nimenrix® (IM)	1	Prophylactic paracetamol	
	Pneumococcal	Prevenar 13® (IM)	3 or 4		
	Meningococcal B	Bexsero® (IM)	3 or 4		
	Hepatitis B	H-B-Vax II® Paed (IM) or Enderix B® Paed (IM)	4		
Hepatitis B	H-B-Vax II® Paed (IM) or Enderix B® Paed (IM)	4	Only for infants born less than 2000g and/or infants born before 32 weeks gestation (regardless of weight)		
18 months	Measles, mumps, rubella, varicella	Priorix-Tetra® (IM) (SC) or ProQuad® (IM) (SC)	2		
	<i>Haemophilus influenzae</i> type b	Act-HIB® (IM)	4		
	Diphtheria, tetanus, pertussis	Infanrix® (IM) or Tripacel® (IM)	4		
	Hepatitis A	VAQTA® (IM)	1		
4 years	Diphtheria, tetanus, pertussis, polio	Infanrix IPV® (IM) or Quadracel® (IM)	5	Maximum 2 doses only See NT Pneumococcal schedule NT Pneumococcal schedule	
	Hepatitis A	VAQTA® (IM)	2		
	Pneumococcal	Pneumovax 23® (IM)	1		
12 years (year 7 students)	Human Papillomavirus HPV	Gardasil 9® (IM)	1	Specified medical risk conditions=3 doses	
	Diphtheria, tetanus, pertussis	Boostrix® (IM) or Adacel® (IM)	1		
14-19 years	Meningococcal ACWY	MenQuadfi® (IM) or Nimenrix® (IM)	2		
	Pneumococcal	Pneumovax 23® (IM)	2	See NT Pneumococcal schedule	NT Pneumococcal schedule

This schedule is a summary. For more detailed information refer to the [The Australian Immunisation Handbook \(health.gov.au\)](https://www.health.gov.au)